




Seagrass-Watch

Proceedings of a workshop for monitoring
seagrass habitats in Queensland

*Northern Fisheries Facility, Cairns
Queensland*

16 - 17 July 2012

Len McKenzie & Rudi Yoshida



First Published 2012

©Seagrass-Watch HQ, 2012

Copyright protects this publication.

Reproduction of this publication for educational or other non-commercial purposes is authorised without prior written permission from the copyright holder provided the source is fully acknowledged.

Reproduction of this publication for resale or other commercial purposes is prohibited without prior written permission of the copyright holder.

Disclaimer

Information contained in this publication is provided as general advice only. For application to specific circumstances, professional advice should be sought.

Seagrass-Watch HQ has taken all reasonable steps to ensure the information contained in this publication is accurate at the time of the survey. Readers should ensure that they make appropriate enquires to determine whether new information is available on the particular subject matter.

The correct citation of this document is

McKenzie, LJ & Yoshida, R.L. (2012). Seagrass-Watch: Proceedings of a workshop for monitoring seagrass habitats in Queensland. Northern Fisheries Facility, Cairns, Queensland, 16 - 17 July 2012 (Seagrass-Watch HQ, Cairns). 50pp.

Produced by Seagrass-Watch HQ

Enquires should be directed to:

Len McKenzie
Seagrass-Watch Program Leader
Northern Fisheries Centre,
PO Box 5396
Cairns, QLD 4870 Australia



Table of Contents

OVERVIEW	5
AGENDA - LEVEL 1 (BASIC)	7
MONDAY 16 TH JULY 2012	7
TUESDAY 17 TH JULY 2012 (GREEN ISLAND)	7
BACKGROUND.....	8
INTERESTING FACTS:.....	16
QUEENSLAND SEAGRASSES. STATUS 2010 - TORRES STRAIT AND EAST COAST	17
GREEN ISLAND SEAGRASSES.....	23
SEAGRASSES OF QUEENSLAND.....	25
PARTS OF A SEAGRASS PLANT.....	27
MONITORING A SEAGRASS MEADOW	29
MONITORING SEAGRASS	30
SEAGRASS-WATCH.....	30
QUARTERLY SAMPLING	33
AT COMPLETION OF MONITORING	34
MAKING A HERBARIUM PRESS SPECIMEN	37
<i>COLLECTION</i>	37
<i>PRESSING</i>	37
<i>HERBARIA</i>	39
UNDERSTANDING SEDIMENT	40
MANAGING SEAGRASS RESOURCES	42
THREATS TO SEAGRASS HABITATS	42
MANAGEMENT	42
REFERENCES	45



Location of Seagrass-Watch long-term monitoring sites, Green Island, Cairns



Overview

Seagrass-Watch is a participatory monitoring program developed in 1998 to provide an early warning of coastal ecological decline. Anyone can participate in Seagrass-Watch, as it responds to local needs. Program participants include scientists, rangers and volunteers (e.g., local residents, indigenous groups, tertiary institutions, non-government organisations). Seagrass-Watch is a monitoring program that brings people together for seagrass conservation. Participants range in ages from 18 to 68 and represent a diverse cross-section of the community, including tradespeople, engineers, school teachers, fishers, divers, retirees, high school and university students, biologists and ecologists. Many are involved with local environmental groups and have a keen interest in conservation and environmental issues.

Seagrass-Watch is a global scientific, non-destructive, seagrass assessment and monitoring program. It identifies areas important for seagrass species diversity and conservation. The information collected can be used to assist the management of coastal environments and to prevent significant areas and species being lost.

Monitoring seagrass resources is important for two reasons: it is a valuable tool for improving management practices; and it allows us to know whether resource status and condition is stable, improving or declining. Successful management of coastal environments (*including seagrass resources*) requires regular monitoring of the status and condition of natural resources.

Early detection of change allows coastal management agencies to adjust their management practices and/or take remedial action sooner for more successful results. Monitoring is important in improving our understanding of seagrass resources and to coastal management agencies for:

- *exposing coastal environmental problems before they become intractable,*
- *developing benchmarks against which performance and effectiveness can be measured,*
- *identifying and prioritising future requirements and initiatives,*
- *determining the effectiveness of management practices being applied,*
- *maintaining consistent records so that comparisons can be made over time,*
- *developing within the community a better understanding of coastal issues,*
- *developing a better understanding of cause and effect in land/catchment management practices,*
- *assisting education and training, and helping to develop links between local communities, schools and government agencies, and*
- *assessing new management practices.*

Seagrass-Watch monitoring efforts are vital to assist with tracking global patterns in seagrass health, and assess the human impacts on seagrass meadows, which have the potential to destroy or degrade these coastal ecosystems and decrease their yield of natural resources. Responsive management based on adequate information will help to prevent any further significant areas and species being lost. To protect the valuable seagrass meadows along our coasts, everyone must work together.

The goals of the Seagrass-Watch program are:

- *to educate the wider community on the importance of seagrass resources*
- *to raise awareness of coastal management issues*
- *to build the capacity of local stakeholders in the use of standardised scientific methodologies*
- *to conduct long-term monitoring of seagrass & coastal habitat condition*
- *to provide an early warning system of coastal environment changes for management*
- *to support conservation measures which ensure the long-term resilience of seagrass ecosystems.*

As part of this Level 1 workshop we will:

- *learn seagrass taxonomy*
- *discuss the present knowledge of seagrass ecology,*
- *discuss the threats to seagrasses*
- *learn techniques for monitoring seagrass resources, and*
- *provide examples of how Seagrass-Watch assists with the management of impacts to seagrass resources and provides an understanding of their status and condition.*

The following information is provided as a training guide and a reference for future Seagrass-Watch monitoring activities. For further information, please do not hesitate to contact us at

Seagrass-Watch HQ

Northern Fisheries Centre

PO Box 5396

Cairns QLD 4870

AUSTRALIA

E-mail hq@seagrasswatch.org

or visit

www.seagrasswatch.org



Photo: Jurgen Freund.

Agenda - Level 1 (*basic*)

Monday 16th July 2012

Morning	0830 - 0845 (15min)	Welcome & Introduction
	0845 - 0900 (15min)	Seagrass Biology and Identification*
	0900 - 0945 (45min)	Classroom activity: seagrass Identification
	0945 - 1030 (45min)	Seagrass Identification <i>continued</i> *
	1030 - 1045 (15min)	<i>Break</i>
	1045 - 1145 (60min)	Seagrass Biology 2 and Ecology
	1145 - 1230 (45min)	Seagrass importance Classroom activity: seagrass importance
Afternoon	1230 - 1315	<i>Lunch</i>
	1315 - 1400 (45min)	Seagrass threats* Classroom activity: seagrass threats
	1400 - 1415 (15min)	Seagrass monitoring*
	1415 - 1515 (60min)	Seagrass-Watch: how to sample* Classroom activity: estimating percent cover
	1515 - 1530 (15min)	<i>Break</i>
	1530 - 1615 (45min)	Seagrass-Watch: how data is used*
	1615 - 1630 (15min)	<i>Wrap-up for day</i>


Tuesday 17th July 2012 (Green Island)

1200 - 1215 (15min)	Safety briefing & risk assessment
1230 - 1415 (105min)	Field exercise: Seagrass-Watch monitoring <i>Where:</i> Green Island (GI1) <ul style="list-style-type: none"> • <i>be punctual</i> • <i>be well rested</i> <i>What to bring:</i> <ul style="list-style-type: none"> • <i>hat, sunscreen (Slip! Slop! Slap!)</i> • <i>dive booties or old shoes that can get wet</i> • <i>drink/refreshments and energising snack</i> • <i>wet weather gear: poncho/raincoat</i> • <i>insect repellent</i> • <i>polaroid sunglasses (not essential)</i> • <i>simple medical kit in case of injuries to yourself</i> • <i>change of footwear and clothes</i> • <i>enthusiasm</i> <p><i>You will be walking across a seagrass meadow along a creek bank exposed with the tide, through shallow water.</i></p>
1415 - 1430	Wrap up (<i>on foreshore</i>) <ul style="list-style-type: none"> • check gear • feedback

Low tide: 1349, 0.5m

Background

Seagrasses are unique flowering plants that have evolved to live in sea water. Seagrasses belong to a group of plants known as angiosperms (flowering plants).



Composite illustration demonstrating morphological features used to distinguish main seagrass taxonomic groups.

Various common names are applied to seagrass species, such as turtle grass, eelgrass, tape grass, spoon grass and shoal grass. These names are not consistently applied among countries. Coastal communities would almost certainly recognise the term “turtle grass” as referring to the shallow subtidal and intertidal seagrasses that turtles are associated with.

Like terrestrial (land living) plants, a seagrass can be divided into its **veins** (lignified conducting tissue that transports food, nutrients and water around the plant), stem, roots (buried in the substrate) and reproductive parts such as flowers and fruits. Algae do not have veins in their leaves nor do they possess roots (anchoring to the surface of the substrate by a holdfast) or produce flowers or seeds.

They are called “seagrass” because most have ribbon-like, grassy leaves. There are many different kinds of seagrasses and some do not look like grass at all. Seagrass range from the size of your fingernail to plants with leaves as long as 7 metres. Some of the shapes and sizes of leaves of different species of seagrass include an oval (paddle or clover) shape, a fern shape, a long spaghetti like leaf and a ribbon shape. Species that have a paddle or fern shaped leaf are called *Halophila*. Ones that have a ribbon shaped leaf are the *Cymodocea*, *Thalassia*, *Thalassodendron*, *Halodule* and *Zostera*. Spaghetti-like seagrass is called *Syringodium*. At the base of a leaf is a sheath, which protects young leaves. At the other end of a leaf is the tip, which can be rounded or pointed. The vertical stem, found in some species, is the upright axis of the plant from which leaves arise (attach). The remnants of leaf attachment are seen as **scars**.

Seagrass leaves lack stomata (microscopic pores on the under side of leaves) but have thin cuticle to allow gas and nutrient exchange. They also possess

Seagrass are marine flowering plants

Seagrasses have roots, stems and leaves

Seagrass is different to seaweed (algae) as they have internal veins, true roots and produce flowers, fruits and seeds

Leaves of different seagrass species can be shaped like a flattened ribbon, look like a fern, round like a clover, or even spaghetti shaped

large thin-walled aerenchyma. The aerenchyma are commonly referred to as veins as they carry water and nutrients throughout the plant (i.e. an internal vascular system). Aerenchyma is specialized tissue having a regular arrangement of air spaces, called lacunae, that both provides buoyancy to the leaves and facilitate gas exchange throughout the plant. Leaves have a very thin cuticle, which allows gas and some nutrient diffusion into them from the surrounding water. Veins can be across the leaf blade or run parallel to the leaf edge. Also within the leaves are chloroplasts, which use the sun's light to convert carbon dioxide and water into oxygen and sugar (photosynthesis). The sugar and oxygen are then available for use by other living organisms.

The roots and horizontal stems (**rhizomes**) of seagrass are often buried in sand or mud. They anchor the plant, store carbohydrates and absorb nutrients. Roots can be simple or branching and all have fine hairs to help absorb nutrients. Rhizomes are formed in segments with leaves or vertical stems rising from the joints, called **nodes** or scars. Sections between the nodes are called internodes. Seagrasses depend upon the growth of rhizomes to increase the area they occupy. This vegetative growth is the most common mode of growth for seagrasses. Although the rhizome mainly runs horizontally, some lateral branches are more or less erect and bear leaves (erect shoots). Sometimes the leaves are on a special type of stalk, called a petiole.

The Roots and Rhizomes of seagrasses are well endowed with aerenchyma and the lacunae are extensive and continuous with leaf tissues. **Oxygen** transport to the roots creates an oxic environment around the roots, facilitating nutrient uptake.

Seagrasses have flowers and pollination systems that are well adapted for pollination by water. Seagrass form tiny flowers, fruits and seeds. Most seagrasses have separate male and female plants. In most species, flowers are small, white and are borne at the base of the leaf clusters. The stamens (male parts) and pistils (female parts) extend above the petals to facilitate pollen release and pollination respectively.

Most seagrasses reproduce by pollination while submerged and complete their entire life cycle underwater. Pollination in seagrasses is hydrophilic (aided by water), and can occur by: (i) pollen transported above water surface (e.g., *Enhalus*); (ii) pollen transported on water surface (e.g., *Halodule*), or; (iii) pollen transported beneath water surface (e.g., *Thalassia*).

Seagrass pollen grains are elongated into a filamentous shape. The filamentous nature of pollen grains facilitates transport within the water medium, mainly by water currents. *Halophila* and *Thalassia* have spherical pollen grains, but they remain joined together in long chains, giving the same effect as having elongated, filamentous pollen grains.

After fertilization, the ovary of the female flower develops into a fruit. In seagrasses, fruit development and fruit structure are as diversified as their flowering patterns and floral structures. In general the seeds, ranging in the size from 0.3 to 0.5mm in some *Halophila* species to more than 1–2 cm in *Enhalus*, are furnished with a nutrition reserve and sink rather than float. The number of seeds within a fruit also varies from 1 (e.g. *Halodule uninervis*) up to 25 (e.g. *Halophila ovalis*).

Seagrass have veins and air channels in their leaves and stems so they can carry water, food and absorb gases

Seagrasses rely on light to convert carbon dioxide and water into oxygen and sugar (photosynthesis)

Roots can be simple or branching and all have fine hairs to help absorb nutrients

Seagrass pump oxygen into the sediment via their roots

Seagrass have flowers, fruits and seeds

Pollination occurs in the water

Pollen from male seagrass flowers is mainly dispersed to female seagrass flowers by tidal currents

Seagrasses are not true grasses

Seagrasses are more closely related to lilies

Seagrass taxonomy

Seagrasses are monocotyledons that are not true grasses (true grasses belong to the family Poaceae), but are rather more closely related to the lily family.

Seagrasses evolved approximately 100 million years ago from land plants that returned to the sea in a least three separate lineages or families. Thus, seagrasses are not a taxonomically unified group but a 'biological' or 'ecological' group. The evolutionary adaptations required for survival in the marine environment have led to convergence (similarity) in morphology.

Worldwide, there are about 12 major divisions, consisting of approximately 60 species (up to 72) of seagrass. The highest concentration of species occurs in the Indo-West Pacific region.

Over 30 species can be found within Australian waters. The most diverse seagrass communities are in the waters of north-eastern Queensland and southern Western Australia.

Various common names are applied to seagrass species, such as turtle grass, eelgrass, tape grass, spoon grass and shoal grass. Seagrasses are not seaweeds. Seaweed is the common name for algae.

Seagrass requirements for growth

Seagrasses require light, nutrients, carbon dioxide, substrate for anchoring, tolerable salinity, temperature and pH to survive. The requirements for a seagrass to be able to exist in the marine environment include:

1. adaptation to life in saline (salty) medium
2. growth when completely submerged
3. anchoring system able to withstand the forces of wave action and tidal currents
4. hydrophilous pollination (pollination aided by water).

The need for physiological adaptations to life in sea water is obvious when one considers that seagrasses evolved from land plants, and most land plants are unable to tolerate even small quantities of salt. In contrast to land plants, some seagrasses can tolerate a salinity range from 4 to 65 parts per thousand (2x seawater concentration). Typically, seagrasses grow best in salinities of 35 parts per thousand. Not all species tolerate all salinities equally well, and salinity tolerance may be a factor promoting different species distributions along salinity gradients, e.g., going up estuaries. Some seagrasses can survive in a range of conditions encompassing fresh water, estuarine, marine, or hypersaline (very salty). A limiting factor for many intertidal seagrasses is osmotic impacts resulting from hypersalinity due to evaporation

Seagrasses being plants need light for photosynthesis. Light availability is the most dominant overriding factor in seagrass growth. Seagrasses have high minimum light requirements (e.g. 10-20% on average, 4.4% minimum and 29% maximum depending on species) of surface irradiance) because: (i) they have a high respiratory demand to support a large non-photosynthetic biomass (e.g. roots, rhizomes); (ii) they lack certain pigments and therefore can utilise only a restricted spectral range; and (iii) they must regularly oxygenate their root zone to compensate for anoxic sediment. However, light in the intertidal can be in excess of requirements and excess light can cause

Seagrass evolved 100 million years ago from land plants that returned to the sea

There are around 60 species of seagrass found in ocean throughout the world

Seagrasses need plenty of sun and clean water to grow.

Seagrasses are physiologically adapted to life in sea water

Seagrasses can tolerate a range of salinities. Some species are less tolerant than others

Light availability is the most important factor determining seagrass growth

Seagrasses require between 10-20% of surface light to grow

temporary photo damage. UV exposure can also have significant impacts on seagrasses.

Temperature influences the rate of growth and the health of plants, particularly at the extremes. As water temperatures increase (up to 38°C) the rate of photorespiration increases reducing the efficiency of photosynthesis at a given CO₂ concentration. The cause of thermal stress at higher temperatures (38°C to 42°C) is the disruption of electron transport activity via inactivation of the oxygen producing enzymes (proteins) of photosystem II. Above these temperatures many proteins are simply destroyed in most plants, resulting in plant death.

Temperature also controls the range of pH and dissolved carbon dioxide (CO₂) concentrations in the water column; factors critical in plant survival in the marine environment.

Seagrasses require inorganic carbon for growth. They uptake inorganic carbon at the leaf surface via two pathways which are species-specific. Some species use bicarbonate (HCO₃⁻) as an inorganic carbon source (e.g. *Halophila ovalis*, *Cymodocea rotundata*, *Syringodium isoetifolium* and *Thalassia*), whereas others use enzymes to make CO₂ available as the inorganic carbon source (e.g. *Enhalus acoroides*, *Halodule*, *Cymodocea serrulata*).

Seagrasses require two key nutrients, nitrogen and phosphorous, for growth. In the coastal regions, seagrasses appear to be primarily limited by nitrogen and secondarily by phosphorus. The demand for nutrients by seagrasses appears to be seasonally dependent. During the growing season the demand for nutrients is high, however during the senescent season elevated nutrients may become toxic.

The availability of nutrients to seagrasses may also be dependent on sediment quality / geochemistry. Bioavailability of nutrients is dependent on particle size and type. For example, clay content influences sediment adsorptive capacity — the more clays the greater the absorptive capacity — and, calcium carbonate binds phosphorus, limiting its bioavailability.

Sediment quality, depth and mobility are important factors for seagrass composition, growth and persistence. Most seagrasses live in sand or mud substrates where their roots and rhizomes anchor the plants to the see floor. Some seagrasses such as *Cymodocea* spp. prefer deeper sediments while others can tolerate a broad range of sediment depths. Colonising seagrasses such as *Halophila* spp. and *Halodule uninervis* are better suited to mobile sediments than larger species. The biogeochemical characteristics of sediment that can affect the nutrient content/binding capacity, organic content and oxygen levels. Seagrasses are unable to grow in sediments of high organic content.

Currents and hydrodynamic processes affect almost all biological, geological and chemical processes in seagrass ecosystems at scales from the smallest (physiological and molecular) to the largest (meadow wide). The pollination of seagrass flowers depends on currents and without current flows, vegetative material and seeds will not be transported to new areas, and species will not be exchanged between meadows. Factors such as the photosynthetic rate of seagrasses depend on the thickness of the diffusive boundary layer that is determined by current flow, as is the sedimentation rate. Both influence growth rates of seagrass, survival of seagrass species and overall meadow morphology.

Water temperature influences the rate of growth and the health of seagrass

Seawater temperatures above 40°C will stress seagrass. Death occurs at temperatures above 43°C

Seagrass require inorganic carbon for growth

Seagrass uptake carbon via two different pathways

Seagrass require two key nutrients, nitrogen and phosphorous, for growth

Nutrient availability to seagrass is dependent on the type of sediment they grow in

Most seagrass live in sand or mud sediments

Sediment movement can determine the presence of seagrass species

Tidal currents are important for pollination and exchange of gases from the water to the plant

Where are seagrasses found?

Seagrasses are found in ocean throughout the world. They occur in tropical (hot), temperate (cool) and the edge of the arctic (freezing) regions. Seagrass are mainly found in bays, estuaries and coastal waters from the mid-intertidal (shallow) region down to depths of 50 or 60 metres. Most species are found in clear shallow inshore areas between mean sea-level and 25 metres depth.

Seagrasses survive in the intertidal zone especially in locations sheltered from wave action or where there is pooling of water at low tide, (e.g., reef platforms and tide pools), which protects seagrass from elevated temperatures and drying.

Seagrasses inhabit all types of ground (substrates), from mud to rock. The most extensive seagrass meadows occur on soft substrates like sand and mud.

The depth range of seagrass is most likely to be controlled at its deepest edge by the availability of light for photosynthesis. Exposure at low tide, wave action and associated turbidity and low salinity from fresh water inflow determines seagrass species survival at the shallow edge.

Seagrass plants form small patches that develop into large continuous meadows. These meadows may consist of one or many species: sometimes up to 12 species present within one location.

How are seagrasses important to the marine ecosystem?

Seagrass communities are one of the most productive and dynamic ecosystems globally. Seagrasses may significantly influence the physical, chemical and biological environments in which they grow by acting as 'ecological engineers'. They provide habitats and nursery grounds for many marine animals and act as substrate stabilisers.

Seagrass meadows are highly productive. They have been documented to create habitat complexity compared with unvegetated areas, providing up to 27 times more habitable substrate, as well as providing refuge and food for a range of animals. About 40 times more animals occur in seagrass meadows than on bare sand.

One of the most important roles of seagrasses is providing a nursery and shelter area for fish and prawns which are valuable to fisheries. Juveniles of some important species which depend on seagrass meadows include fish such as perch, mullet, whiting, tailor, bream, snappers, emperors and sweetlips. Commercial penaeid prawns such as red spot king, brown tiger, grooved tiger and endeavour also live in seagrass meadows as juveniles. Tropical rock lobsters also live in seagrass meadows as juveniles. Shellfish such as some oysters and pearl shell may be more likely to settle and survive where there is seagrass. Juvenile and adult sandcrabs and flathead are just two species which spend most of their lives in seagrass meadows, where there is not only food but also protection from strong tidal currents and predators. Larger predatory animals such as herons, cormorants, sharks, barramundi, salmon, crocodiles, etc, are also attracted to the seagrass meadows by the schools of forage fish which seek shelter there.

Seagrass meadows are a major food source for a number of grazing animals and are considered very productive pastures of the sea. The dugong

Seagrass are commonly found in estuaries, shallow coastal locations, and on reef-tops.

Seagrass are mainly found in clear shallow inshore areas between mean sea-level and 25 metres depth.

The depth that seagrass are found underwater depends on the light availability (water clarity)

Seagrass plants form small patches that develop into large meadows

Seagrasses are important habitat and feeding grounds for marine organisms.

About 40 times more animals occur in seagrass meadows than on bare sand.

Seagrasses are important nursery grounds for fish, and they support many human commercial activities.

(*Dugong dugon*) and the green turtle (*Chelonia mydas*) mainly feed on seagrass. An adult green turtle eats about two kilograms of seagrass a day while an adult dugong eats about 28 to 40 kilograms a day. Although dugongs and turtles will feed on any seagrass species within their range, if a range of species is available, they select seagrass species for food which are high nitrogen, high starch and low fibre. For example, the order of seagrass species preference for dugongs is *Halophila ovalis* > *Halodule uninervis* > *Zostera capricorni*. In sub-tropical and temperate areas, water birds such as black swans also eat seagrass.

Decomposing seagrasses provide food for benthic (bottom-dwelling) aquatic life. The decaying leaves are broken down by fungi and bacteria which in turn provide food for other microorganisms such as flagellates and plankton. Microorganisms provide food for the juveniles of many species of marine animals such as fish, crabs, prawns and molluscs.

The rhizomes and roots of the grasses bind sediments on the substrate, where nutrients are recycled by microorganisms back into the marine ecosystem. The leaves of the grasses slow water flow, allowing suspended material to settle on the bottom. This increases the amount of light reaching the seagrass meadow and creates a calm habitat for many species.

Seagrasses are nutrient sinks, buffering or filtering nutrient and chemical inputs to the marine environment. Seagrasses uptake nitrogen and phosphorus from coastal run-off that, in overabundance, can lead to algal blooms that can impair water quality.

Interactions with mangroves and coral reefs

Tropical seagrasses are important in their interactions with mangroves and coral reefs. All these systems exert a stabilizing effect on the environment, resulting in important physical and biological support for the other communities).

Barrier reefs protect coastlines, and the lagoon formed between the reef and the mainland is protected from waves, allowing mangrove and seagrass communities to develop. Seagrasses trap sediment and slow water movement, causing suspended sediment to fall out. This trapping of sediment benefits coral by reducing sediment loads in the water.

Mangroves trap sediment from the land, reducing the chance of seagrasses and corals being smothered. Sediment banks accumulated by seagrasses may eventually form substrate that can be colonized by mangroves. All three communities trap and hold nutrients from being dispersed and lost into the surrounding oceanic waters.

Valuation of seagrasses

The valuation of ecosystem services is a very controversial topic in today's literature. Ecosystem Services are the processes by which the environment produces resources that we often take for granted. For seagrasses it is services such as clean water, preventing erosion, and habitat for fisheries. The economic values of seagrass meadows are very large, although not always easy to quantify. Seagrass meadows are rated the 3rd most valuable ecosystem globally (on a per hectare basis), only preceded by estuaries and wetlands. The average global value of seagrasses for their nutrient cycling services and the raw product they provide has been estimated at 1994 US\$ 19,004 ha⁻¹ yr⁻¹.

Dugongs can eat up to 40kg of seagrass per day.

Dugongs and turtles select seagrass species for food which are high nitrogen, high starch and low fibre

Seagrasses also contribute to the productivity of ecosystems via the detrital food pathway

Seagrass binds sediments and help prevent erosion

Seagrasses slow water flow and increase water clarity

Seagrass help remove harmful nutrient and sediment pollution from coastal waters

Seagrasses, mangroves and coral reef interact, providing physical and biological support for other communities

Seagrass meadows are rated the 3rd most valuable ecosystem globally (more valuable than mangroves or coral reefs)

What causes seagrass areas to change?

Tropical seagrass meadows vary seasonally and between years, and the potential for widespread seagrass loss has been well documented.

Factors which affect the distribution of seagrass meadows are sunlight and nutrient levels, water depth, turbidity, salinity, temperature, current and wave action.

Seagrasses respond to natural variations in light availability, nutrient and trace element (iron) availability, grazing pressure, disease, weather patterns, and episodic floods and cyclones. The dynamic nature of seagrass meadows in response to natural environmental variation complicates the identification of changes caused by humans.

What threatens seagrass?

Seagrass meadows can be easily damaged. Approximately 58% of seagrass meadows globally, have lost part of their distribution. According to reports, the documented losses in seagrass meadows globally since 1980 are equivalent to two football fields per hour.

Some losses are natural due to storms and herbivores, however most losses are the result of human activities. Human pollution has contributed most to seagrass declines around the world.

The most widespread and pervasive cause of seagrass decline is a reduction in available light. Processes that reduce light penetration to seagrasses include pulsed turbidity events during floods, enhanced suspended sediment loads and elevated nutrient concentrations. Poor farming practices can result in excess sediments and fertilizers washing down creeks to the sea. Sewage discharge and stormwater runoff from urban development can elevate nutrients in coastal areas. Boating activity may also stir up sediment, reducing light levels. Phytoplankton and fast-growing macroalgae are also better competitors for light than benthic plants and their biomass can shade seagrasses during progressive eutrophication.

Oil and trace metal contamination can exert direct toxic effects on some seagrass species. Seagrasses are able to bioaccumulate the trace metals and this can have ramifications for grazers such as dugongs.

People can also physically damage or destroy seagrass. Coastal development for boat marinas, shipping ports and housing generally occurs on the coast in areas which are sheltered and seagrass like to grow. Seagrass meadows are either removed or buried by these activities. Coastal developments can also cause changes in water movement. Dredging boat channels to provide access to these developments not only physically removes plants, but can make the water muddy and dump sediment on seagrass. Litter and rubbish can also wash into the sea if not properly disposed. Rubbish can physically and chemically damage seagrass meadows and the animals that live within them.

Boating and fishing activities can physically impact or destroy seagrasses. Boat anchors and their chains can dig into seagrass. Propellers can cut into seagrass meadows and unbalance the rhizome mat. Storms can further exacerbate the damage by the physical force of waves and currents ripping up large sections of the rhizome mat. Uncontrolled digging for bait worm can also physically damage seagrasses and some introduced marine pests and pathogens also have the potential to damage seagrass meadows.

Seagrasses can change due to both natural and human impacts

People can damage or destroy seagrass by pollution (sewage, oil spills and coastal runoff) and physical destruction (dredging, bait & clam digging, boat propellers and anchors/moorings).


Poor farming practices can result in runoff which can damage seagrass by elevating nutrients, reducing available light and releasing herbicides.

Coastal development can have a major impact on seagrass meadows

One of the other significant impacts to seagrass is climate change. The major vulnerability of seagrass to climate change is loss of seagrass in the coastal zone, particularly near river mouths and in shallow areas. The greatest impact is expected to result from elevated temperatures, particularly in shallower habitats where seagrasses grow (e.g., effecting distribution and reproduction). In addition, reduced light penetration from sediment deposition and resuspension are expected due to more intensive cyclones/hurricanes and elevated flooding frequency and amplitude. This will result in even greater seagrass losses, and changes in species composition are expected to occur particularly in relation to disturbance and recolonisation. Following such events, a shift to more ephemeral species and those with lower minimum light requirements is expected.

Climate change can threaten intertidal seagrass by increased seawater temperature and greater physical disturbance from storms

Please note: citations have been removed for ease of reading. Please see *References & Further Reading* for source/citations on scientific facts.



Notes:

.....

.....

.....

.....

.....

.....

.....



Interesting facts:

Over a billion people live within 50 km of a seagrass meadow. Millions of people obtain their protein from animals that live in seagrasses.

The estimated global coverage of seagrass is between 300,000 and 600,000 square kilometres.

A hectare of seagrass absorbs 1.2 kilograms of nutrients per year, equivalent to the treated effluent from 200 people.

In northern Australia, whole seagrass meadows are able to completely replace their leaves (turnover) in around 14 days during the growing season.

A hectare of seagrass sequesters 830 kilograms of carbon per year, equivalent to the CO₂ emissions from an automobile travelling 3,350 km.

One square metre of seagrass can produce up to 10 litres of oxygen per day

In northern Australia, the primary productivity of seagrass meadows is higher than a mangrove forest, a terrestrial forest or grassland.

Seagrasses occupy only 0.1% of the seafloor, yet are responsible for 12% of the organic carbon buried in the ocean, which helps reduce greenhouse gases.

The only endangered marine plant is a species of seagrass (*Halophila johnsonii* in Florida).

There is a single clone of seagrass that is over 6,000 years old (*Posidonia oceanica* in the Mediterranean Sea). It is possibly the world's oldest plant!

The deepest growing seagrass (*Halophila decipiens*), 86 metres, was reported from Cargados Carajos Shoals in the Indian Ocean northeast of Mauritius.

Seagrass produce the longest pollen grains on the planet.

Some intertidal species of seagrasses can lose up to 50% per cent of their water content and still survive.

Did you know that Australia has the highest number of seagrass species of any continent in the world?


In Alaska, seagrasses remain frozen and in a dormant state over winter and do not start to grow again until the thaw.

The longest known seagrass 7.3 metres in length has been reported from Funakoshi Bay, Japan.

40,000 seeds of *Halodule uninervis* have been found in 1 square metre of mudflat.

In Florida, 80% of the above ground seagrass biomass is consumed by parrot fish.

The anchor and chain from one cruise boat can destroy an area of seagrass the size of a football field!



Queensland seagrasses Status 2010 – Torres Strait and East Coast

Seagrasses are a key component of the marine ecosystem of Queensland and are essential for sustainable and productive fisheries.

There are 15 seagrass species common in Queensland waters, a region that is part of the Indo Pacific centre of seagrass biodiversity. Seagrasses are widespread and found from the upper intertidal region, on reef platforms, and down to 70 metres below sea level. They are one of the key habitat types along with algae and coral that underpin the biodiversity and productivity of tropical ecosystems and their fisheries. The Torres Strait and the east coast of Queensland have some of the most extensive seagrass meadows in the world with an estimated 17,206 sq km in

the strait and at least 38,079 sq km down the east coast. As the status of the seagrass meadows and risks from human influences is not consistent over this large area, we have summarised our data by Natural Resource Management (NRM) areas. The threats to seagrass are almost all in the southern half of Queensland (south of Cooktown). Seagrass meadows in Queensland have declined in abundance down the east coast over the last 3-4 years, however the total area of seagrass in Queensland has changed little over the long-term (5-10 years). Seagrass declines where they have occurred are most likely the result of natural variations in climate, particularly tropical storms and flood run-off, against a background of reduced water quality.



Biology of seagrass

Seagrasses are flowering plants that have evolved to live in sea water. They are found in shallow inshore areas of bays, estuaries and coastal waters throughout the world. Fifteen of the approximately 70 seagrass species known, are common in Queensland waters.


Seagrass provides a nursery and shelter for fish, prawns and shellfish. Seagrass meadows have up to 27 times more habitable surface and support about 40 times more animals compared to bare seafloor. Seagrass is the primary food source for dugong, endangered green turtles, fish and numerous other animals that live in, on or around seagrasses. In some parts of the world, the carbon sink capacity of seagrass ecosystems is nearly equivalent to that of undisturbed rainforest.

It is estimated that the global value of ecosystem services that seagrass meadows provide is around US\$1.9 trillion per year in nutrient cycling alone.

Seagrasses are also important because they reduce coastal erosion and filter toxic chemicals and nutrient run-off. The health of seagrass meadows is an excellent indicator of the health of inshore waters and early detection of change allows government agencies to adjust their management response.

Seagrass distribution

Approximately 18,374 square kilometres of seagrass has been mapped in the coastal waters around Queensland and Torres Strait since the mid 1980s. Surveys and statistical modelling of seagrass in offshore waters deeper than 15m shows 37,454 square kilometres of the sea floor within the Great Barrier Reef World Heritage Area and Torres Strait has some seagrass present making Queensland's seagrass resources globally significant.



General conceptual model of seagrass habitats in north east Australia (from Carruthers et al. 2002)

Seagrass habitats

The diversity of seagrass species in Queensland reflects the variety of habitats provided by coasts, bays, estuaries, and reefs as well as the complexity of the Torres Strait island systems and the length of the Great Barrier Reef and inshore lagoon. Seagrass habitats can be separated into four major categories: estuary/inlet, coastal, reef and deepwater. All but the outer reef habitats are influenced by seasonal pulses of sediment-laden, nutrient-rich river flows, resulting from high volume summer rainfall. Cyclones, severe storms, wind and waves as well as macro grazers (fish, dugongs and turtles) influence all habitats in this region. The result is dynamic, spatially and temporally variable seagrass meadows.



Cymodocea rotundata



Cymodocea serrulata



Enhalus acoroides



Halodule pinifolia



Halodule uninervis

Threats & risks

Over the last few decades the area of seagrass around the world has been declining at a rate of two football fields per hour. This places seagrass meadows among the most threatened ecosystems. Fortunately, threats and impacts in Queensland are low in most locations compared to other countries and the area of seagrass in Queensland has remained relatively stable.

Agricultural run-off poses the largest human-induced threat to seagrass health. Seagrass is also threatened, in decreasing order of magnitude, by the following impacts: urban and industrial runoff; urban and port infrastructure development; dredging; shipping accidents; trawling; recreational boat damage; commercial boat damage and netting.

As a result of successful, cooperative fisheries management in Queensland, fishing poses a relatively low threat to seagrass health.

Bottom trawling is restricted by spatial and temporal closures and trawl operators avoid areas of known seagrass meadows. A recent reduction in commercial boat numbers has also reduced pressure on bottom habitats.

Relatively small human populations along much of the coast and the successful management of key threats through legislation such as the *Fisheries Act 1994* provide some respite to implement strategies to ensure the long-term survival of seagrass resources. Fisheries Queensland has a responsibility to manage and safeguard these vital habitats. Seagrass resource status information is one tool to enable successful management.

Map of risks to seagrass

Coastal areas that are dark blue have a high likelihood of seagrass presence and are areas where the risk of impacts from human activities is low.



Coastal areas that are red have a high likelihood of seagrass presence and are areas where the risk of impacts from human activities is high.

Seagrass in Queensland exposed to a high cumulative risk profile are all in ports or close to areas of urban development such as Cairns, Townsville, Mackay and Gladstone.

The risk map is for the Great Barrier Reef World Heritage Area only.

In Queensland, all marine plants, including seagrass, are protected from unlawful damage. Incidental damage and limited collection of seagrass is allowed only in accordance with the DEEDI self assessable code MP05 or by permit (see www.dpi.qld.gov.au/28_9109.htm).

0 100 200
kilometres



Halophila capricorni



Halophila decipiens











Halophila minor



Halophila ovalis



Halophila spinulosa



Legend: Seagrass maps are composite (all mapping surveys pooled). Actual meadow boundary 50% - 100% Modelled meadow probabilities (pixel size of 2km²) from all deepwater (>15m) survey points pooled. Detailed maps & sources are available at www.dpi.qld.gov.au/28_142.htm



Halophila tricostata



Syringodium isoetifolium




Thalassia hemprichii



Thalassodendron ciliatum



Zostera muelleri subsp. *capricorni*



Relative seagrass abundance (all sites pooled) for intertidal estuarine, coastal and reef habitats within each NRM region along the east coast of Queensland (pooled by season for each year). Long-term trend line is calculated as polynomial. Values in parenthesis are the number of monitoring units (sites or meadows). Seagrass abundance (seagrass percentage cover or biomass as g DW m⁻²) relative to the 95th percentile at each monitoring site enables standardised comparisons.

The data underpinning this assessment of seagrass status along the east-coast of Queensland and Torres Strait has been collected from the Seagrass-Watch program, with additional input from the Habitats at Risk and the Risk Modelling programs. The Seagrass-Watch program is a participatory monitoring program involving community groups and agencies. Seagrass-Watch plays a key role in tracking changes to coastal seagrass meadows in Queensland, nationally and globally, providing early warning of coastal ecological decline.

The Habitats at Risk program is a partnership with industry to provide expert advice on coastal, fishing and port management to ensure Queensland develops its industries with the least impact to seagrass habitats. The Risk Modelling program using available data sets to provide advice on management options and to assess the ecosystem values of seagrass meadows.

NRM	Area of seagrass (km ²)	# species	mean % cover	abundance data quality	status (since Jan2007)
Torres Strait	<15m = 13,413 >15m = 3,793	total = 13 estuary = 2 coastal = 11 reef = 11	58.4 ±1.2 (Feb 2007-Mar 2010) 36.6 ±0.5 (Jun 2004-Mar 2010)	deficient inadequate sufficient	
Cape York	<15m = 1,843 >15m = 10,641	total = 15 estuary = 4 coastal = 6 reef = 9	17.5 ±0.5 (Oct 2003-Apr 2010)	deficient deficient inadequate	
Wet Tropics	<15m = 201 >15m = 6,638	total = 14 estuary = 5 coastal = 5 reef = 9	22.6 ±0.7 [^] (Dec 1993-Nov 2009) 12.4 ±0.2 (Aug 2000-Apr 2010) 30.3 ±0.4 (Nov 2001-Apr 2010)	inadequate sufficient sufficient	
Burdekin	<15m = 55 >15m = 2,372	total = 11 estuary = 3 coastal = 8 reef = 8	11.3 ±1.4 (Sep 2001-Jun 2004) 19.6 ±0.3 (Apr 2001-Jun 2010) 30.3 ±0.6 (May 2005-Jun 2010)	deficient sufficient adequate	
Mackay Whitsunday	<15m = 154 >15m = 293	total = 12 estuary = 4 coastal = 5 reef = 8	20.7 ±0.4 (Jun 2000-Mar 2010) 20.1 ±0.2 (Sep 1999-Mar 2010) 12.0 ±0.2 (Apr 2000-Mar 2010)	inadequate sufficient adequate	
Fitzroy	<15m = 240 >15m = 6,725	total = 6 estuary = 4 coastal = 3 reef = 4	18.0 ±0.5 (Oct 2005-Apr 2010) 19.8 ±0.2 (Apr 2003-Apr 2010) 2.3 ±0.2 (Sep 2007-Apr 2010)	sufficient adequate inadequate	
Burnett Mary	<15m = 1,026 >15m = 6,992	total = 8 estuary = 8 coastal = 5	9.9 ±0.1 (Aug 1999-May 2010) 3.8 ±0.1 (Aug 1999-Feb 2010)	sufficient adequate	
SEQ	<15m = 401 >15m = unknown	total = 9 estuary = 8 coastal = 7	35.0 ±0.2 (May 2001-Dec 2008) 33.7 ±0.5 (Feb 2003-Jul 2008)	sufficient sufficient	

Area calculated from composite (all mapping surveys pooled) maps. Area measures for seagrass >15m depth determined by actual mapping and modelled probabilities 50% (pixel size of 5km²) from all survey points pooled. NB: actual and modelled information is at different levels of precision.

[^]Abundance in blue = grams Dry Weight m² above-ground. Abundance data quality (column 5) indicates level of precision: deficient = no or very few monitoring sites; inadequate = several sites but not enough to be representative; adequate = enough sites to be representative; sufficient = many sites available.

Status: increasing or stable decreasing variable/uncertain unknown

Future

In the next 10-20 years it is expected that changes in climate (i.e. increasing temperature, sea-level rise and tropical storm frequency) will increase stress to coastal habitats. This will add to existing threats to seagrass meadows. The best defence against unnecessary habitat loss will be to use high quality, long-term information in decision making regarding coastal development and management. Monitoring change in seagrass performance and understanding the factors underpinning

trends in change, resilience and recovery processes will be critical. There are significant gaps in our knowledge of seagrass ecology, namely: species level response to change, subtidal meadow dynamics, responses to light quality and turbidity, and in understanding responses to natural cycles such as La Niña and El Niño fluctuations. The Marine Ecology Group is working in partnership with James Cook University and other research institutions to improve knowledge in these areas.

We thank all Queensland Seagrass-Watch participants whose dedication and data collection over the past decade have made this brochure possible. We thank M. Rasheed, H. Taylor, M. Waycott (JCU) and the Marine Ecology Group for their contributions and also the many organisations who assisted with monitoring in partnership with Fisheries Queensland.

Fisheries Queensland has a team of scientists in the Marine Ecology Group that conduct state, national and global seagrass assessment, monitoring and research. The group provides the scientific papers, reports and advice that in partnership with managers, industry and other research organisations underpins the successful management, conservation and protection of fisheries habitat.

Further reading

Carruthers et al. (2001). Seagrass habitats of north east Australia: models of key processes and controls. *Bulletin of Marine Science* 71(3): 1153-1169.

Coles et al. (2007). *Status and trends of seagrass habitats in the Great Barrier Reef World Heritage Area*. RRRC Ltd, Cairns (122 pp). www.rrrc.org.au

Coles et al. (2003). The seagrasses of eastern Australia. *World Atlas of Seagrasses*, eds. Green & Short. UNEP WCMC, UCLA Press, Berkeley USA.

Larkum et al. (2006). *Seagrasses: Biology Ecology and Conservation*. Springer, The Netherlands. 691 pp.

Waycott et al. (2007). Vulnerability of seagrasses in the Great Barrier Reef to climate change. *Climate Change and the Great Barrier Reef*, eds. Johnson & Marshall. GBRMPA and Australian Greenhouse Office, Australia. pp.193-235.

This brochure should be cited as: L. McKenzie, R. Yoshida, A. Grech & R. Coles (2010) Queensland seagrasses. Status 2010 – Torres Strait and East Coast. Fisheries Queensland (DEEDI), Cairns. 6pp

Why conserve seagrass?



Seagrasses are **economically and ecologically** valuable to both humans and marine life. Seagrass is one of the most productive natural ecosystems in the world.

Seagrasses **improve water quality** by acting as nutrient sinks, buffering or filtering nutrient and chemical inputs to the marine environment. They also **stabilise** coastal sediments, helping to avert erosion.

Seagrasses provide **food and shelter** for many organisms (shrimps, crabs, worms, snails and small fish) and are a nursery ground for **commercially important** prawn and fish species.

Larger fish and seabirds visit seagrass meadows to feed. Endangered sea turtles and dugongs also graze on seagrass.

Seagrass meadows are fragile ecosystems. Human impacts such as excessive pollution from sewage discharge, oil spills, herbicides, coastal runoff, dredging, boat propellers and anchors/moorings can damage or destroy seagrasses.



How you can help

In Queensland, all marine plants, including seagrass, are protected from unlawful damage. Incidental damage and limited collection of seagrass is allowed if only in accordance with the DPI&F self assessable code MPO5 (see website below).

Many efforts are underway to educate the public about the benefits of seagrass and how they can help to protect seagrass. There are many ways you can help: don't litter; be aware when applying fertilizers and pesticides, as excess amounts can wash down gutters and drains to the sea; when boating, slow down and avoid shallow areas; support marine conservation initiatives; learn about these special marine habitats and volunteer to monitor their health by joining Seagrass-Watch.

Seagrass-Watch is a global seagrass assessment and monitoring program. Seagrass-Watch monitoring efforts are vital to assist with tracking global patterns in seagrass health, and assess the human impacts which have the potential to destroy or degrade these coastal ecosystems and decrease their yield of natural resources. Responsive management based on adequate information will help to prevent any further significant areas and species being lost. To protect the valuable seagrass meadows along our coasts, everyone must work together.



For more information visit:
www.seagrasswatch.org

Seagrasses of Green Island

Cymodocea rotundata

- flat, strap-like leaves 2-4mm wide
- rounded, smooth leaf tip
- smooth rhizome
- leaf scars form continuous ring around the stem
- found on shallow reef flats



Cymodocea serrulata

- strap-like leaves, 5-9mm wide
- leaf tip serrated
- leaf sheath is broadly triangular
- leaf scars not continuous ring around the stem
- found on shallow subtidal reef flats



Halodule uninervis

- trident leaf tip
- 1 central longitudinal vein
- rhizome usually pale with clean black leaf scars
- dugong & turtle preferred food



Halophila capricornii

- small oval leaves that are hairy on one side
- central vein on leaf with 9-14 cross veins
- usually found > 10m depth in coral environments
- proximal to coral reefs



Halophila decipiens

- small oval leaf blade 1-2.5cm long
- 6-8 cross veins
- leaf hairs on both sides
- leaves usually longer than wider
- found at subtidal depths (>10m)



Halophila ovalis

- oval shaped leaves in pairs
- 8 or more cross veins
- smooth leaf surface
- preferred dugong food



Halophila spinulosa

- fern like
- leaves arranged in opposite pairs
- erect shoot up to 15cm long
- found at subtidal depths (>3m)



Syringodium isoetifolium

- narrow, cylindrical spaghetti-like leaves
- leaves 7-30cm long, taper to a point
- 2-3 leaves arising at each shoot
- rhizomes thin



Thalassia hemprichii

- broad ribbon like, curved leaves
- short black bars of tannin cells in leaf blade
- thick rhizome with scars between shoots
- common on reef flats



Seagrasses of Green Island



Seagrass-Watch

www.seagrasswatch.org

Seagrass-Watch HQ
Northern Fisheries Centre
Po Box. 5396, Cairns QLD 4870
AUSTRALIA
Email: hq@seagrasswatch.org
Phone: +61 7 4057 3731
www.seagrasswatch.org



Text and design by L. McKenzie and R. Yoshida, seagrass waterloobers by G. Berry.

What are Seagrasses?



Seagrasses are unique marine flowering plants of which there are approximately 60 species worldwide. Various common names are applied to seagrass species, such as turtle grass, eelgrass, tape grass, spoon grass and shoal grass. Seagrasses are not seaweeds. Seaweed is the common name for algae.

Seagrass live in sheltered coastal waters, undergo pollination while submerged and complete their entire life cycle underwater. They grow much like land grasses, with extensive below ground rhizomes or runners. Plants form small patches that develop into large continuous meadows. These meadows may consist of one or many species, sometimes up to 12 species present within one location.


Because seagrass requires sunlight, most seagrass is found in the intertidal zone especially in locations sheltered from wave action or where there is pooling of water at low tide, (e.g., reef platforms and tide pools), which protects seagrass from elevated temperatures and drying.



Seagrasses of North Qld



Seagrasses are a major component of north Queensland's wet tropics marine ecosystems. Thirteen species of seagrass have been recorded, representing 87% of the known species found in Queensland waters. The wide range of physical habitats where seagrasses are found undoubtedly contributes to the high species diversity. Habitats include intertidal and subtidal areas of estuary, coastal, reef and deepwater environments.



General conceptual model of seagrass habitats in north east Australia

Seagrasses of Green Island

Green Island is a vegetated coral cay located approximately 27 km north-east of Cairns. Nine species of seagrasses are commonly found on the reef flat and sea bed surrounding Green Island.

Scattered over the reef flat you will find **Cymodocea rotundata** and the hooked leaved **Thalassia hemprichii**. On the edge of the meadows adjacent to the beach, you will find **Halodule uninervis** with its three pointed leaf tips and the clover like **Halophila ovalis**, both species are food for dugong. If you look closely on the shallow reef near the beach here, you'll find some broad leaved **Cymodocea serrulata** with its serrated leaf tips. In the shallow lagoon out from the jetty, you'll find **Syringodium isoetifolium** with it's distinctive spaghetti-like leaves and a mixture of broader leaved **Halodule uninervis** and **Cymodocea rotundata**. These are heavily cropped by green turtles which feed around the island.


In the deeper (>3m) sandy areas off the reef edge, you can find hairy clover like **Halophila decipiens** and the fern like **Halophila spinulosa**, both favoured food for dugong. To the north west of the reef, you may even find the rare **Halophila capricorni**, a clover like species which has hairs on only one side of its leaves.

Seagrasses on Green Island are being monitored as part of the Great Barrier Reef Water Quality Protection Plan because they are important indicators of marine environment 'health'. The monitoring programme provides a critical component of the assessment of any long-term improvement in water quality that will occur as best land management practices are widely adopted across the Great Barrier Reef catchments and regions.



HC

HD HS



HD

HS

HO

HU SI

HU HO SI

CS HU SI TH

CR CS HU SI TH

SI


HU HO

CR HU

HU HO

CR HU TH

CR TH



CR

CS

HU

HO

SI

TH

TH

Seagrasses of Queensland

Adapted from Waycott, M, McMahon, K, Mellors, J., Calladine, A., and Kleine, D (2004) A guide to tropical seagrasses in the Indo-West Pacific. (James Cook University Townsville) 72pp.

Leaves cylindrical



cylindrical

Syringodium isoetifolium

- leaves taper to a point
- leaves contain air cavities
- inflorescence a “cyme”
- leaves 7-30cm long

Ruppia maritima


- leaves fine and thread-like
- pointed tip on leaves, sometimes serrated
- leaves up to 15cm long
- rhizome fragile
- inflorescence on a long stalk, sometimes spiralled

Leaves oval to oblong



oval to oblong

obvious vertical stem with more than 2 leaves




Halophila spinulosa

- leaves arranged opposite in pairs
- leaf margin serrated
- shoots can be up to 15cm long
- 10-20 pairs of leaves per shoot
- leaf 15-20mm long and 3-5mm wide

Halophila tricostata

- leaves arranged in clusters of 3, at a node on vertical stem
- leaf margin sparsely serrated
- leaf clusters do not lie flat
- 5-12 leaf clusters per shoot
- leaf 12-20mm long and 2-4mm wide

leaves with petioles, in pairs



Halophila capricorni

- leaf margins finely serrated
- fine hairs on one side of leaf blade
- leaf 15-30mm long and 5-9 mm wide
- 9-14 cross vein pairs, occasionally forked

Halophila decipiens

- leaf margins finely serrated
- fine hairs on both sides of leaf blade
- leaf apex rounded to slightly pointed
- leaf 10–25mm long and 3–10mm wide
- 6-8 cross vein pairs

Halophila minor

- less than 8 pairs of cross veins
- leaf 5-15mm long and 3.5-6mm wide
- leaf margins smooth
- no leaf hairs

Halophila ovalis


- cross veins 8 or more pairs
- leaf 5-40mm long and 5-20mm wide
- leaf margins smooth
- no leaf hairs

Leaves strap-like

Leaves can arise from vertical stem



straplike



Cymodocea rotundata

- leaf tip rounded with smooth edge
- leaf 2-4mm wide with 9-15 parallel veins
- leaf sheath scars continuous around stem
- old sheaths forming a fibrous mass at the base of each shoot

Cymodocea serrulata

- leaf tip rounded with serrated edge
- leaf 4-9mm wide with 13-17 parallel veins
- leaf sheath broadly flat and triangular, not fibrous
- leaf sheath scars not continuous around upright stem

Halodule uninervis

- leaf tip tri-dentate or pointed, not rounded
- leaf blades 0.5-5mm wide
- leaf with 3 distinct parallel veins, sheaths fibrous
- rhizome usually white with small black fibres at the nodes

Halodule pinifolia

- leaf tip rounded
- narrow leaf blades 0.25-1.2mm wide
- leaf with 3 distinct parallel veins, sheaths fibrous
- rhizome usually white with small black fibres at the nodes


Thalassia hemprichii

- leaf tip rounded, may be slightly serrated
- leaf 4-12mm wide with 9-11 parallel veins
- leaf with obvious red flecks, 1-2mm long
- leaf often distinctly curved
- rhizome thick with distinct scars, usually triangular in shape
- one short root per rhizome node

Thalassodendron ciliatum

- distinct upright stem
- clusters of curved leaves (>5 mm wide), margins serrated
- stem and rhizome woody

Leaves always arise directly from rhizome




Enhalus acoroides






- large plant, leaves >30 cm long, >1 cm wide
- in-rolled edges of leaves
- long, black bristles protruding from thick rhizome

Zostera muelleri subsp. *capricorni*

- leaf with 3-5 parallel-veins
- cross-veins form boxes
- leaf tip smooth and rounded, may be dark point
- rhizome usually brown or yellow in younger parts
- prophyllum present, i.e. single leaf originating from rhizome instead of from vertical, leaf bearing shoot.


Parts of a seagrass plant



Leaf		
Tip	Can be rounded or pointed. Tips are easily damaged or cropped, so young leaves are best to observe.	 <p style="text-align: center;"><i>rounded</i> <i>pointed</i></p>
Veins	Used by the plant to transport water, nutrients and photosynthetic products. The pattern, direction and placement of veins in the leaf blade are used for identification. <ul style="list-style-type: none"> • cross-vein: perpendicular to the length of the leaf • parallel-vein: along the length of the leaf • mid-vein: prominent central vein • Intramarginal-vein: around inside edge of leaf 	 <p style="text-align: center;"><i>cross</i> <i>parallel</i></p> <p style="text-align: center;"><i>mid</i> <i>intramarginal</i></p>
Edges	The edges of the leaf can be either serrated, smooth or inrolled	 <p style="text-align: center;"><i>serrated</i> <i>smooth</i> <i>inrolled</i></p>
Sheath	A modification of the leaf base that protects the newly developing tissue. The sheath can entirely circle the vertical stem or rhizome (continuous) or not (non-continuous); fully or partly cover the developing leaves and be flattened or rounded. Once the leaf has died, persistent sheaths may remain as fibres or bristles.	 <p style="text-align: center;"><i>clean & flattened</i> <i>fibrous</i></p>
Attachment	The leaf can attach directly to the rhizome, where the base of the leaf clasps the rhizome, or from a vertical stem or stalk (petiole) e.g. <i>Halophila ovalis</i> .	 <p style="text-align: center;"><i>petiole</i> <i>scar</i> <i>stem</i></p> <p style="text-align: center;"><i>rhizome</i></p>


Stem

The vertical stem, found in some species, is the upright axis of the plant from which leaves arise (attach). The remnants of leaf attachment are seen as scars. Scars can be closed (*entirely circle the vertical stem*) or open (*do not entirely circle the vertical stem*).




Rhizome

The horizontal axis of the seagrass plant, usually in sediment. It is formed in segments, with leaves or vertical stem arising from the joins of the segments, the nodes. Sections between the nodes are called internodes. Rhizomes can be fragile, thick and starchy or feel almost woody and may have scars where leaves were attached.



Root

Underground tissues that grow from the node, important for nutrient uptake and stabilisation of plants. The size and thickness of roots and presence of root hairs (very fine projections) are used for identification. Some roots are simple or cordlike, others may be branching, depending on seagrass species.



Notes:

.....

.....

.....

.....

.....



Monitoring a seagrass meadow

Monitoring is the repeated observation of a system, usually to detect change. It is an integrated activity to evaluate the condition of the physical, chemical and biological character of the environment. Environment monitoring programs provide coastal managers with information and assist them to make decisions with greater confidence.

Environmental monitoring programs are ideally designed to: quantify the causes of change; examine and assess acceptable ranges of change for the particular site; and to measure levels of impacts.

Common drivers (reasons) for monitoring include: community interest; government policies such as Coastal Strategies and Plans, Oceans Policy, State of the Environment Reporting (SoE), Water Quality guidelines or Best Practice Guidelines; and Government Legislation (e.g., Fish Habitat Protection).

Users of the monitoring program information/results are diverse, including for example: the general public, environmental regulators - legislators, resource managers and scientists.

There are a number of issues to consider when implementing a monitoring program, including: ensure the protocols used have explicit objectives; clearly identified responsibilities of the partners (e.g. Gov agencies, consultants, community groups); a clear and defensible rationale for using the parameters that are measures (e.g. physico/chemico, biological indicators); to have a baseline (first) assessment / measure against which subsequent changes can be measured/compared; knowledge of spatial and temporal variation prior to designing the program (i.e. pilot study); clearly defined field protocols; data management procedures, ensure the level of change and accuracy of the detection is appropriate (as will vary according to the methodology); selection of statistical tools; and a mechanism to reduce and manage errors (i.e. QA/QC program).

Appropriate Quality Assurance/Quality Control (QA/QC) procedures are an integral component of all aspects of sample collection and analysis in monitoring programs. This includes participation in relevant inter-laboratory studies, proficiency testing, and the use of standard reference materials. Monitoring programs often include the following guidelines for implementation by data collectors and reporters:

- appropriate methods must be in place to ensure consistency in field procedures to produce robust, repeatable and comparable results including consideration of sampling locations, replication and frequency;
- all methods used must be fit for purpose and suited to a range of conditions;
- appropriate accreditation of participating laboratories or provision of standard laboratory protocols to demonstrate that appropriate laboratory QA/QC procedures are in place for sample handling and analysis;
- participation in inter-laboratory performance testing trials and regular exchange of replicate samples between laboratories;
- rigorous procedures to ensure 'chain of custody' and tracking of samples;
- appropriate standards and procedures for data management and storage; and
- a process to ensure data collectors are aware of any errors and provide an opportunity to clarify or correct data.



Monitoring seagrass

Seagrasses are often at the downstream end of catchments, receiving runoff from a range of agricultural, urban and industrial land-uses. Seagrass communities are generally susceptible to changes in water quality and environmental quality that make them a useful indicator of environmental health. Seagrass make good **bioindicators** of environmental health because they are:

- are widely distributed;
- have an important ecological role;
- are sessile plants which show measurable and timely responses to external stressors/impacts (rather than relocating to a less stressful environment) and;
- are integrative of environmental conditions.

Several factors are important for the persistence of healthy seagrass meadows, these include: sediment quality and depth; water quality (temperature, salinity, clarity); current and hydrodynamic processes; and species interactions (e.g., epiphytes and grazers). Seagrass generally respond in a typical manner that allows them to be measured and monitored. In reporting on the health of seagrasses it is important to consider the type of factors that can effect growth and survival. Factors include:

- increased turbidity reduces light penetration through the water, interfering with photosynthesis and limiting the depth range of seagrass;
- increased nutrient loads encourages algal blooms and epiphytic algae to grow to a point where it smothers or shade seagrasses, thereby reducing photosynthetic capacity;
- increased sedimentation can smother seagrass or interferes with photosynthesis;
- herbicides can kill seagrass and some chemicals (e.g., pesticides) can kill associated macrofauna;
- boating activity (propellers, mooring, anchors) can physically damage seagrass meadows, from shredding leaves to complete removal;
- storms, floods and wave action can rip out patches of seagrasses.

Seagrass-Watch

A simple method for monitoring seagrass resources is used in the Seagrass-Watch program. This method uses standardised measurements taken from sites established within representative intertidal meadows to monitor seagrass condition. The number and position of sites can be used to investigate natural and anthropogenic impacts.

Seagrass-Watch is one of the largest seagrass monitoring programs in the world. Since it's genesis in March 1998 in Australia, Seagrass-Watch has now expanded internationally to more than 26 countries. Monitoring is currently occurring at over 300 sites. To learn more about the program, visit www.seagrasswatch.org.

Seagrass-Watch aims to raise awareness on the condition and trend of nearshore seagrass ecosystems and provide an early warning of major coastal environment changes. Participants of Seagrass-Watch are generally volunteers from a wide variety of backgrounds who all share the common interest in marine conservation. Most participants are associated with established local community groups, schools, universities & research institutions, government (local & state) or non-government organisations.

Seagrass-Watch integrates with existing education, government, non-government and scientific programs to raise community awareness to protect this important marine habitat for the benefit of the community. The program has a strong scientific underpinning with an



emphasis on consistent data collection, recording and reporting. Seagrass-Watch identifies areas important for seagrass species diversity and conservation and the information collected is used to assist the management of coastal environments and to prevent significant areas and species being lost.

Seagrass-Watch methods were developed to be rigorous, yet relatively simple and easy to use. Each of the parameters used have been carefully chosen with a clear and defensible rationale. The protocols used have explicit objectives and the sampling strategy is prepared using baseline and knowledge of spatial and temporal variation. As the collection of data by a diversity of participants necessitates a high level of training to ensure that the data is of a standard that can be used by management agencies. After 6–9 hours of training, participants can produce reliable data. Training includes both formal and informal approaches. Formal training is conducted by Seagrass-Watch HQ for participants over 17 years of age and includes formal lectures and on-site assessments with a tiered level of certification for competency. Formally trained participants are certified to supervise on-site monitoring and demonstrate (i.e. informally train) monitoring methods. Informal training is also conducted by local coordinators and/or scientists. Ideally, at least one formally trained volunteer is present at each monitoring event. Evidence of competency is securely filed at Seagrass-Watch HQ.

Seagrass-Watch has an accepted Quality Assurance-Quality Control program in place to ensure that the program is producing data of high quality, and that time and resources are not wasted. Seagrass-Watch HQ has systems in place to manage the way Seagrass-Watch data is collected, organised, documented, evaluated and secured. The Seagrass-Watch program collects and collates all data in a standard format. By using simple and easy methods, Seagrass Watch ensures completeness (the comparison between the amounts of valid or useable data originally planned to collect, versus how much was collected). Standard seagrass cover **calibration sheets** are used to ensure precision (the degree of agreement among repeated measurements of the same characteristic at the same place and the same time) and consistency between observers and across sites at monitoring times to ensure percentage covers are close to a true or standardised value.

Other QAQC procedures include the selection of intertidal seagrass sites which are **permanently marked** with either plastic star pickets or an accurate (± 3 m) GPS waypoint. Labels identifying the sites and contact details for the program are attached to these pickets. Positions of 0 m and 50 m points for all three transects at a site are also noted using GPS. This ensures that the same site is monitored each event and that data can be compared between periods of time.

Ongoing standardisation of observers is achieved by on-site refreshers of standard percentage covers by all observers prior to monitoring and through ad hoc comparisons of data returned from duplicate surveys (e.g. either a site or a transect will be repeated by scientist – preferably the next day and unknown to volunteers). Any discrepancy in these duplicates is used to identify and subsequently mitigate bias. For the most part however uncertainties in percentage cover or species identification are mitigated in the field via direct communication (as at least one experienced/certified observer is always present), or the collection of voucher specimens (to be checked under microscope and pressed in herbarium) and the use of a digital camera to record images (protocol requires at least 27% of quadrats are photographed) for later identification and discussion.

Seagrass-Watch HQ has implemented a quality assurance management system to ensure that data collected is organised and stored and able to be used easily. All data (datasheets and photographs) received are entered onto a relational database on a secure server. Receipt of all original data hardcopies is documented and filed within the Queensland Government Registered Management System, a formally organised and secure system. Seagrass-Watch HQ operates as custodian of data collected from other participants and provides an evaluation and



analysis of the data for reporting purposes. Access to the IT system and databases is restricted to only authorised personnel. Provision of data to a third party is only on consent of the data owner/principal.

Seagrass-Watch HQ checks all data for completeness, consistency and accuracy. All data submitted to Seagrass-Watch HQ it is first checked from compliancy:

- *legible original datasheets,*
- *good quality quadrat photographs (high resolution),*
- *voucher specimens (if required) and*
- *completed MS Excel spreadsheet.*

Validation is provided by checking observations against photographic records to ensure consistency of observers and by identification of voucher specimens submitted. In accordance with QA/QC protocols, Seagrass-Watch HQ advises observers via an official **Data Error Notification** of any errors encountered/identified and provides an opportunity for correction/clarification (this may include additional training).

Once Seagrass-Watch HQ has completed all checks, a field in the Master database identifies data as either passed, quarantined, non-compliant or not-passed. Non-compliant data is used for large-scale summary reporting only if the data quality is deemed acceptable, i.e. if it was collected by a Level 1 trained participant, that the scans/copies of datasheets are OK (*only if originals are not available*), and/or that the quadrat images were acceptable to complete QAQC, etc. If data quality is unacceptable, the data is either not entered into the Master database or remains quarantined/not-passed (excluded from analysis & reporting). If predominantly non-compliant data is used for detailed analysis and reporting at a site or location/region, it is marked on the outputs with a notice of non-compliance (e.g., site graphs). If officially requested data is non-compliant, a note in the metadata advises of non-compliance and includes a caveat to "use with caution". Any data considered unsuitable (e.g. nil response to data notification within thirty days) is quarantined or removed from the database.

Seagrass-Watch employs a proactive approach to monitoring, involving ongoing training for participants and the continued development of new methods and refinement of existing methods, including location/habitat specific calibration sheets, operation & validation of autonomous temperature and light loggers, etc. Quality data reassures the data users (e.g., coastal management agencies) that they can use the data to make informed decisions with confidence.

Seagrass-Watch monitoring efforts are vital to assist with tracking global patterns in seagrass health, and assessing human impacts on seagrass meadows, which have the potential to destroy or degrade these coastal ecosystems and decrease their value as a natural resource. Responsive management based on adequate information will help to prevent any further significant areas and species being lost. To protect the valuable seagrass meadows along our coasts, the community, government and researchers have to work together.


THE GOALS OF THE PROGRAM ARE:

- *To educate the wider community on the importance of seagrass resources*
- *To raise awareness of coastal management issues*
- *To build the capacity of local stakeholders in the use of standardised scientific methodologies*
- *To conduct long-term monitoring of seagrass & coastal habitat condition*
- *To provide an early warning system of coastal environment changes for management*
- *To support conservation measures which ensure the long-term resilience of seagrass ecosystems.*

Seagrass-Watch Protocols

Source: McKenzie, L.J., Campbell, S.J., Vidler, K.E. & Mellors, J.E. (2007) *Seagrass-Watch: Manual for Mapping & Monitoring Seagrass Resources*. (Seagrass-Watch HQ, Cairns) 114pp (www.seagrasswatch.org/manuals.html)

Site layout



Quadrat code = site + transect+quadrat

e.g., P11225 = Pigeon Is. site 1, transect 2, 25m quadrat

Pre-monitoring preparation

Make a Timetable

Create a timetable of times of departure and arrival back, and what the objective of the day is and what is to be achieved on the day. Give a copy of this to all volunteers involved in advance so they can make their arrangements to get to the site on time. List on this timetable what the volunteers need to bring.

Have a Contact Person

Arrange to have a reliable contact person to raise the alert if you and the team are not back at a specified or reasonable time.

Safety

- Assess the risks before monitoring - check weather, tides, time of day, etc.
- Use your instincts - if you do not feel safe then abandon sampling.
- Do not put yourself or others at risk.
- Wear appropriate clothing and footwear.
- Be sun-smart.
- Adult supervision is required if children are involved
- Be aware of dangerous marine animals.
- Have a first aid kit on site or nearby
- Take a mobile phone or marine radio

Necessary equipment and materials

- 3x 50metre fibreglass measuring tapes
- 6x 50cm plastic tent pegs
- Compass
- 1x standard (50cm x 50cm) quadrat
- Magnifying glass
- 3x Monitoring datasheets
- Clipboard, pencils & 30 cm ruler
- Camera & film
- Quadrat photo labeller
- Percent cover standard sheet
- Seagrass identification sheets

Quarterly sampling

Within the 50m by 50m site, lay out the three 50 transects parallel to each other, 25m apart and perpendicular to shore (see site layout). Within each of the quadrats placed for sampling, complete the following steps:

Step 1. Take a Photograph of the quadrat

- Photographs are usually taken at the 5m, 25m and 45m quadrats along each transect, or of quadrats of particular interest. First place the photo quadrat labeller beside the quadrat and tape measure with the correct code on it.
- Take the photograph from an angle as **vertical** as possible, which includes the entire quadrat frame, quadrat label and tape measure. Avoid having any shadows or patches of reflection off any water in the field of view. Check the photo taken box on datasheet for quadrat.

Step 2. Describe sediment composition

- Dig your fingers into the top centimetre of the substrate and feel the texture. Describe the sediment by noting the grain size in order of dominance (e.g., Sand, Fine sand, Fine sand/Mud).

Step 3. Describe other features and ID/count of macrofauna

- Note and count (whole numbers - never use < or > symbols) any features which may be of interest (e.g. gastropods, hermit crabs, turtle feeding) within the comments column.

Step 4. Estimate seagrass percent cover

- Looking down on the quadrat from above, estimate the total percentage of the seabed (substrate) within the quadrat covered by seagrass. Estimate the footprint/shadow provided by the seagrass shoots.
- Always use the percent cover photo standards (calibration sheets) as your guide, estimating cover as accurate as possible, e.g. 27%, 61%
- If cover is below 3%, you can count the seagrass shoots and calculate percent cover using the rule of 1 shoot = 0.1%

Step 5. Estimate seagrass species composition

- Identify the species of seagrass within the quadrat and determine the percent contribution of each species (starting with least abundant, total composition must equal 100%)
- Use seagrass species identification keys provided and use more than 1 feature to identify the species

Step 6. Measure seagrass canopy height

- Measure canopy height (in centimetres) of the dominant strap-leaf species, ignoring the tallest 20%.
- Measure from the sediment to the leaf tip of 3 shoots, entering all 3 measures onto datasheet

Step 7. Estimate algae percent cover

- Looking down on the quadrat from above, estimate the total percentage of the seabed (substrate) within the quadrat covered by macroalgae (independent of seagrass cover)
- Macroalgae is not attached to seagrass leaves and may be attached to rocks, shells or may be drift

Step 8. Estimate epiphyte percent cover

- Epiphytes are algae attached to seagrass blades and often give the blade a furry appearance.
- First estimate how much of an average seagrass leaf surface is covered, and then how many of the leaves in the quadrat are covered. For example, if 20% of the blades are each 50% covered by epiphytes, then quadrat epiphyte cover is 10%. Use the epiphyte matrix to assist you.
- Do not include epifauna with epiphytes. Epifauna are sessile animals attached to seagrass blades – record % cover of epifauna in the comments or an unused/blank column – do not add to epiphyte cover.

Step 9. Take a voucher seagrass specimen if required

- Place seagrass samples in a labelled plastic bag with a little seawater and a waterproof label. Select a representative specimen of the species and ensure that you have all the plant parts including the rhizomes and roots. Collect plants with fruits and flowers structures if possible.

Step 10. Move to next quadrat

- Repeat steps 1 to 8 for the remaining 32 quadrats

Step 11. At completion of monitoring

- Check data sheets are filled in fully.
- Remove equipment from site (eg non-permanent pegs)

At completion of monitoring

Step 1. Wash & pack gear

- Rinse all tapes, pegs and quadrats with freshwater and let them dry.
- Review supplies for next quarterly sampling and request new materials
- Store gear for next quarterly sampling

Step 2. Press any voucher seagrass specimens if collected

- The voucher specimen should be pressed as soon as possible after collection. Do not refrigerate longer than 2 days, press the sample as soon as possible.
- Allow to dry in a dry/warm/dark place for a minimum of two weeks. For best results, replace the newspaper after 2-3 days.

Step 3. Submit all data

- Data can be entered into the MS-Excel file downloadable from www.seagrasswatch.org. Email completed files to hq@seagrasswatch.org
- Mail original datasheets, photos and herbarium sheets


Seagrass-Watch HQ
Northern Fisheries Centre
PO Box 5396
Cairns QLD 4870 AUSTRALIA

SEAGRASS SPECIES CODES

Ho

Halophila ovalis

- 8 or more cross veins
- no hairs on leaf surface
- leaf margins smooth
- leaf 5-20mm long



Halophila decipiens

- small oval leaf blade 1-2.5cm long
- 6-8 cross veins
- leaf hairs on both sides
- found at subtidal depths




Hd

Hm

Halophila minor

- less than 8 pairs of cross veins
- small oval leaf blade less than 5mm wide
- leaf margins smooth
- no leaf hairs



Hu

Halodule uninervis

- flat leaf, 0.25-5mm wide
- trident leaf tip
- 1 central vein
- usually pale rhizome, with clean black leaf scars



Cs

Cymodocea serrulata

- serrated leaf tip
- wide leaf blade (5-9mm wide)
- leaves 6-15cm long
- 13-17 longitudinal veins
- robust/strong rhizome



Hs

Halophila spinulosa

- fern like
- leaves arranged in opposite pairs
- leaves flat & serrated
- erect shoot to 15cm long
- found at subtidal depths



Si

Syringodium isoetifolium

- narrow spaghetti-like leaves
- cylindrical in cross section, 1-2mm diameter
- leaves contain air cavities
- leaf tip tapers to a point
- leaves 7-30cm long
- fleshy white rhizomes



Zc

Zostera muelleri subsp. *capricorni*

- flat leaf, 2-5mm wide
- leaf with 3-5 parallel-veins
- cross-veins form boxes
- leaf tip smooth and rounded, may be dark point at tip
- leaf grows directly from rhizome ie no stem
- rhizome usually brown or yellow in younger parts





SEAGRASS-WATCH MONITORING



ONE OF THESE SHEETS IS TO BE FILLED OUT FOR EACH TRANSECT YOU SURVEY

START of transect (GPS reading)

Latitude: 25° 11.2818' S Longitude: 152° 37.5372' E

OBSERVER: Bev Citizen DATE: 17 / 2 / 09
 LOCATION: Burrum Heads
 SITE code: BH1 TRANSECT no.: 2
 START TIME: 1304 END TIME: 1340

Quadrat (metres from transect origin)	Sediment (eg. mud/sand/shell)	Comments (eg 10x gastropods, 4x crab holes, dugong feeding trails, herbarium specimen taken)	Seagrass coverage (%)	% Seagrass species composition				Canopy height (cm)	% Algae cover	% Epi- cover
				HO	HU	ZC	Water			
1 (0m)	Sand	SC x 3 HC x 1	40	30	70		0	514.7	5	33
2 (5m)	S	GAB x 3	33	50	50		0	1017.8	10	18
3 (10m)	mud/sand	worm x 1	18	70	20	10	0	618.5	0	48
4 (15m)	m s	DFT x 1	0				0	0	17	0
5 (20m)	m s shell	HC x 3	36	5	90	5	1cm	917.5	12	57
6 (25m)	m s sh	Turtle cropping	48	100			1cm	NA.	2	96
7 (30m)	Fine Sand		0				1.5cm	0	23	0
8 (35m)	FS	SC x 2 CH x 3	0.7		100		2cm	717.7	18	31
9 (40m)	S m		23	96	4		2cm	214.6	6	17
10 (45m)	m	Mudworek x 2 HC x 1	41	2	95	3	2cm	5.5, 6 9	3	21
11 (50m)	m s		16	3	7	90	2cm	716.7	38	6

END of transect (GPS reading)

Latitude: 25° 11.2656' S Longitude: 152° 37.5546' E

SC = Sea Cucumbers
 GAB = Gastropod
 DFT = Dugong feeding trail
 HC = Hermit Crab
 CH = Crab Hole



Making a herbarium press specimen

Herbaria are repositories of preserved and labelled plant specimens, arranged to allow easy access and archival storage. The specimens are typically in the form of herbarium sheets: pressed and dried plants that have been attached to a sheet of heavy paper together with a data label. A herbarium specimen is simple in form and low-tech in preparation, yet it preserves a wealth of valuable information. If properly stored, a herbarium specimen will last for centuries without much deterioration. Specimens document the variation in form and geographical range of species. Herbaria also document valuable historical collections, such as "type specimens", the original specimens on which a plant's Latin name is based. Many herbarium specimens record the existence of plants in habitats now developed and lost.

COLLECTION

Before collecting any seagrass specimens, ensure you have the appropriate permits.

In the field, collect a handful of representative seagrass shoots, including the leaves, rhizomes and roots. Keep in mind that it is not always possible to get a successful classification if you do not have particular parts such as flowers, fruits, seeds and roots, so try to select shoots which have these features. Ideally, collect plants with growing tips (meristems) as they contain higher concentrations of DNA which could aid genetic identification in the future.

Specimens should be pressed as soon as possible after collection. If it is more than 2 hours before you press the specimen, then you should refrigerate to prevent any decomposition. Do not refrigerate longer than 2 days, press the sample as soon as possible.

PRESSING

Tools

First you will need some clean white cartridge-type paper (photocopy paper will suffice) and herbarium sheets (if available). You will also need forceps, scissors/scalpel, a dish of clean fresh water and a herbarium press. It is not difficult to build a home-made press, keeping in mind that what must be accomplished is to keep the specimens squeezed between layers of paper (newspapers or blotting paper) until they are totally devoid of the original content of water. The upper and lower parts of the press might be made of heavy cardboard or thick plywood or equivalent material. A more advanced kind of press might be built for an optimal drying of your plants. This press can be made with two wooden boards with screws and nuts placed at each corner: turning the nuts the two boards will come closer pushing together the paper with the plants. This kind of press can be built at home or bought in some art tools stores.

Preparation

Wash the seagrass specimen in clean fresh water and carefully remove any debris, epiphytes or sediment particles.

Arrangement

It is very important that the seagrass specimen be arranged so that you can immediately see all the main characters of that particular species; so do not focus only at the aesthetics of the mounted specimen. It is advisable to arrange specimens before being placed in the press as once dried, plant specimens can easily be broken if handled without care. The best manner to place the plants on the mounting sheets is to align them with the right side of the page (or diagonally if space is required) and to have the heaviest parts and specimens at the bottom. Leaves can be folded in larger specimens if a larger press is not available. It is better to leave an empty space at the borders of the mounting sheets; but you can either arrange your specimens (along with the label) in a regular way from page to page, or stagger the specimens

at different positions on each sheet, so that each group of sheets will have a more equally distributed pressure.

Labels

Each specimen must have a label on its own sheet, which should include the taxonomic denomination (*at least family, genus and species*) along with information on the date and place of collection. The name of the collector and of the individual who did the determination should also be added. Use permanent and water resistant ink (black or blue) to write your labels; otherwise a pencil can be used (medium lead). Specimen labels should include:

- species name (*if known*)
- location & site code (*if applicable*)
- date collected
- latitude/longitude
- water depth
- % seagrass cover
- sediment type
- other seagrass species present
- name of collector and who identified the specimen
- comments -*such as presence of flowers/fruits or ecological notes*

Place the label on the lower right hand corner of the paper.

Drying

Place another clean sheet of paper over the specimen and place within several sheets of newspaper. As circulating air is very important to get your specimens dried in a short time, the assemblage of specimen/paper should be placed within two sheets of corrugated cardboard and then into a herbarium press. Corrugated cardboard ensures air can penetrate and speed up the drying process. If no corrugated cardboard is available, keep the filled press size small.

Once in the herbarium press, wind down the screws until tight (*do not over tighten*). If you do not have a press, the specimens can be pressed by putting some heavy object on top, i.e. bricks or large books. It is important that the plants are put under sufficient pressure; otherwise more time will be required to achieve a good desiccation, besides they could be damaged by dampness and moulds.



The press should be exposed to a gentle heat source, avoiding excessive heat that will "cook" the specimens. Sometimes it is possible to use the heat from the sun. In this case the presses should be small. If fire is the heat source, keep the press at a safe distance to prevent fire starting on the press.

Changing the paper is a very important step. In the first three or four days a paper change should take place every day, then you can leave more time between changes. If you neglect the change of paper the plants will take more time to lose their water content, besides they could be damaged if the paper stays wet for a few days. When changing the paper you must keep the specimens intact and ensure the label travels with the specimen. The minimum time required for complete drying ranges from two to four days or more. Once a specimen has become dry and stiff, it can be mounted and placed into the herbarium.

Mounting

Once the specimen is completely dry, you will need to mount it to herbarium sheets if available or a new clean white cartridge-type paper.

There are different ways to mount the specimens to the herbarium sheets, such as strapping, gluing, pinning or nothing. We recommend the strapping method using removable adhesive tape (e.g. Magic Tape). The tape pulls off easily, leaves behind no messy residue, and can be pulled up and moved around. To fix the specimen to the mounting paper, lay small strips of tape across a few sturdy parts of the plant (e.g. either end of rhizome or a stem) at a minimal number of points. This method will allow a certain degree of movement for further examinations, but the specimen will not fall from the mounting paper

HERBARIA

Once the specimen is mounted it can be stored in a dry place or lodged in Herbaria. If you do not have a Herbaria in your region or state (usually located at a University or Government agency), you can submit specimens to Seagrass-Watch HQ which maintains a Herbaria as part of the Australia Tropical Herbarium.

Alternatively, you can email a scanned image of the pressed specimen. Please ensure that the scanned image is no less than 600 dpi and includes the specimen and label. Scanned images can be sent to hq@seagrasswatch.org and will be lodged in the Seagrass-Watch Virtual Herbarium <http://www.seagrasswatch.org/herbarium.html>.

The Virtual Herbarium is an electronic gateway to the collections of the Seagrass-Watch HQ herbaria. The goals of the Virtual Herbarium are to make specimen data available electronically for use in biodiversity research projects; to reduce transport of actual specimens for projects where digital representations will suffice for study; and to provide a source of reference information for Seagrass-Watch participants.

Understanding sediment

Seagrasses, especially structurally large species, affect coastal and reef water quality by trapping sediments and acting as a buffer between catchment inputs and reef communities. Seagrass meadows have the ability to modify the energy regimes of their environments, and help stabilise sediment by trapping and binding the sediment. However, the trapping ability of seagrass is in reality an equilibrium established between deposition/sedimentation and erosion/resuspension.

Studies have shown that sediment characteristics are important in determining seagrass growth, germination, survival, and distribution. As part of Seagrass-Watch, field descriptions of sediment type collected 0-2 cm below the sediment/water interface are determined by visual and tactile inspection of (wet) samples and constituents (primary descriptors) differentiated according to the Udden – Wentworth grade scale.

Grain size classes used, based on the Udden – Wentworth grade scale of Wentworth (1922).

	Fine-medium Clay	0 – 0.002 mm
	Coarse Clay	0.0021 – 0.004 mm
Mud	Very Fine Silt	0.0041– 0.008 mm
	Fine Silt	0.0081 – 0.016 mm
	Medium Silt	0.0161 – 0.031 mm
	Coarse Silt	0.0311 – 0.063 mm
	<hr/>	
	Very Fine Sand	0.0631 – 0.125 mm
	Fine Sand	0.1251 – 0.250 mm
Sand	Medium Sand	0.2501 – 0.500 mm
	Coarse Sand	0.5001 – 1.000 mm
	Very Coarse Sand	1.0001 – 2.000 mm
<hr/>		
Gravel	Granules	2.0001 – 4.000 mm
	Pebbles and larger	>4.0001 mm


In Seagrass-Watch, the primary descriptors relate to the size of the sediment grains: gravel (>2000µm); coarse sand (>500 µm); sand (>250 µm); fine sand (>63 µm); and mud (<63 µm).

The sediment **Primary Descriptors** are written down from left to right in decreasing order of abundance: e.g. Mud/Sand is mud with sand, where mud is determined as the dominant constituent (by volume).


mud	<i>has a smooth and sticky texture.</i>
fine sand	<i>fairly smooth texture with some roughness just detectable. Not sticky in nature.</i>
sand	<i>rough grainy texture, particles clearly distinguishable.</i>
coarse sand	<i>coarse texture, particles loose.</i>
gravel	<i>very coarse texture, with some small stones.</i>

Sediment type **Modifiers** are also commonly used, however these are recorded in the comments section. Modifiers include: coral, shell grit, forams, diatoms, etc.


The visual/tactile estimation method used in Seagrass-Watch is a simple yet relatively accurate measure of the sediment grain size which can be used for quantitative assessments (see McKenzie 2007, http://www.seagrasswatch.org/Info_centre/Publications/pdf/371_DPIF_McKenzie.pdf).




gravel (>2mm)




very coarse sand (1 - 2 mm)




coarse sand (0.5 - 1 mm)



medium sand (0.25 - 0.5 mm)



fine sand (0.125 - 0.25 mm)



very fine sand (0.063 - 0.125mm)

Notes:

.....

.....

.....

.....

.....



Managing seagrass resources

Threats to seagrass habitats

Destruction or loss of seagrasses have been reported from most parts of the world, often from natural causes, e.g., "wasting disease" or high energy storms. However, destruction commonly has resulted from human activities, e.g., as a consequence of eutrophication or land clamation and changes in land use. Increases in dredging and landfill, construction on the shoreline, commercial overexploitation of coastal resources, and recreational boating activities along with anthropogenic nutrient and sediment loading has dramatically reduced seagrass distribution in some parts of the world. Anthropogenic impacts on seagrass meadows continue to destroy or degrade coastal ecosystems and decrease the function and value of seagrass meadows including their contribution to fisheries. It is possible global climate change will have a major impact. Efforts are being made toward rehabilitation of seagrass habitat in some parts of the world: transplantation, improvement of water quality, restrictions on boating activity, fishing and aquaculture, and protection of existing habitat through law and environmental policy.

Management

Seagrasses do not exist in nature as a separate ecological component from other marine plants and are often closely linked to other community types. In the tropics the associations are likely to be complex interactions with mangrove communities and coral reef systems. In temperate waters, algae beds, salt marshes, bivalve reefs, and epiphytic plant communities are closely associated with areas of seagrass. Many management actions to protect seagrasses have their genesis in the protection of wider ecological systems or are designed to protect the overall biodiversity of the marine environment.

Seagrasses are also food for several marine mammal species and turtles, some of which (such as the dugong *Dugong dugon* and green turtle *Chelonia mydas*) are listed as threatened or vulnerable to extinction in the IUCN Red List (www.iucnredlist.org). Seagrasses are habitat for juvenile fish and crustaceans that in many parts of the world form the basis of economically valuable subsistence and/or commercial fisheries. The need to manage fisheries in a sustainable way has itself become a motivating factor for the protection of seagrasses.

Methods of direct protection range from legislative instruments and associated legal sanctions through to education (Coles & Fortes 2001). These can be separated into three approaches: a proscriptive legal approach; a non-proscriptive broad based approach ranging from planning processes to education; and a reactive approach designed to respond to a specific issue such as a development proposal. These may overlap and be used simultaneously in many cases. It is these three approaches that Seagrass-Watch supports for the protection/conservation of seagrass.

Reactive (on-ground)

Reactive processes generally occur in response to a perceived operational threat such as a coastal development proposal (Coles & Fortes 2001). Reactive processes can include port contingency planning, risk management plans and environmental impact assessments.

Prescriptive (legal)

Prescriptive management of seagrass issues can range from local laws to a Presidential Decree, or Executive Order. Laws can directly safeguard seagrasses



or can protect them indirectly by protecting habitat types (all aquatic vegetation) or by influencing a process, e.g., prevention of pollution (Coles & Fortes 2001).

In some locations, protection is often strongest at the village or community level. This may be by Government supported agreements or through local management marine area level. In these cases successful enforcement is dependent on community support for the measure.

Non-prescriptive (planning & education)

Non-prescriptive methods of protecting seagrasses are usually part of planning processes and may have a strong extension/education focus (Coles & Fortes 2001). Providing information is important as it enables individuals to voluntarily act in ways that reduce impacts to seagrasses. Non-prescriptive methods range from simple explanatory guides to complex industry codes of practice.

Coastal management decision making is complex, and much of the information on approaches and methods exists only in policy and legal documents that are not readily available. There may also be local or regional Government authorities having control over smaller jurisdictions with other regulations and policies that may apply. Many parts of South East Asia and the Pacific Island nations have complex issues of land ownership and coastal sea rights. These are sometimes overlaid partially by arrangements put in place by colonising powers during and after World War II, leaving the nature and strength of protective arrangements open for debate.

Both Australia and the United States have developed historically as Federations of States with the result that coastal issues can fall under State or Federal legislation depending on the issue or its extent. In contrast, in Europe and much of South East Asia, central Governments are more involved. Intercountry agreements in these areas such as the UNEP Strategic Action Plan for the South China Sea and the Mediterranean Countries Barcelona Convention (<http://www.unep.org/>) are required to manage marine issues that encompass more than one country.

Approaches to protecting seagrass tend to be location specific or at least nation specific (there is no international legislation directly for seagrasses as such that we know of) and depend to a large extent on the tools available in law and in the cultural approach of the community. There is, however, a global acceptance through international conventions (RAMSAR Convention; the Convention on Migratory Species of Wild Animals; and the Convention on Biodiversity) of the need for a set of standardised data/information on the location and values of seagrasses on which to base arguments for universal and more consistent seagrass protection.

Indigenous concepts of management of the sea differ significantly from the introduced European view of the sea as common domain, open to all and managed by governments (Hardin 1968). Unlike contemporary European systems of management, indigenous systems do not include jurisdictional boundaries between land and sea. Indigenous systems have a form of customary ownership of maritime areas that has been operating in place for thousand of years to protect and manage places and species that are of importance to their societies.

Marine resource management these days should, therefore, attempt to achieve the following interrelated objectives: a) monitor the wellbeing (e.g. distribution, health and sustainability) of culturally significant species and environments (e.g. dugong, marine turtles, fish, molluscs, seagrass etc.); and b) monitor the cultural values associated with these culturally significant species and environments (Smyth *et al.* 2006).



To realize objective a) we believe the following also needs to be accomplished if the successful management of coastal seagrasses is to be achieved.

1. Important fish habitat is known and mapped
2. Habitat monitoring is occurring
3. Adjacent catchment/watershed impacts and other threats are managed
4. Some level of public goodwill/support is present
5. Legal powers exist that are robust to challenge
6. There is effective enforcement and punishment if damage occurs

The key element is a knowledge base of the seagrass resource that needs to be protected and how stable/variable that resource is. It is also important to know if possible any areas that are of special value to the ecosystems that support coastal fisheries and inshore productivity. It is important as well that this information is readily available to decision makers in Governments in a form that can be easily understood.

Consequently a combination of modern “western” science and indigenous knowledge should be brought together within a co-management framework for the successful management of these resources. (Johannes 2002; Aswani & Weiant 2004; Turnbull 2004; Middlebrook and Williamson 2006; Gaskell 2003, George et al. 2004). This can only occur if the resource owners actively involve themselves in the management of their resources. Western science also needs to recognise that resource owners have practical and spiritual connections with the resources found within their environment. Once this is recognized then this approach will have the added benefit of empowering communities who own the knowledge to be the primary managers and leaders in decisions about their land and sea country.



References

- Aswani, S., Weiant, P. (2004). Scientific evaluation in women's participatory management: monitoring marine invertebrate refugia in the Solomon Islands. *Human Organisation* **63** (3), 301-319.
- Coles R.G. and Fortes, M. (2001) Protecting seagrass - approaches and methods. In Short, F.T. and Coles, R.G. (eds) 2001. *Global Seagrass Research Methods*. Elsevier Science B.V., Amsterdam. Chapter 23, pp. 446-463.
- Gaskell, J. (2003). Engaging science education within diverse cultures. *Curriculum Inquiry*. **33**: 235-249.
- George, M., Innes, J., Ross, H. (2004). Managing sea country together: key issues for developing co-operative management for the Great Barrier Reef World Heritage Area. CRC Reef Research Centre Technical Report No 50, CRC Reef Research Centre Ltd, Townsville.
- Hardin, G. (1968). The tragedy of the commons. *Science, New Series* **162** (3859), 1243-1248.
- Johannes, R.E. (2002). The renaissance of community-based marine resource management in Oceania. *Annu. Rev. Ecol. Syst.* **33**: 317-340.
- McKenzie, L.J. (2007) Relationships between seagrass communities and sediment properties along the Queensland coast. Progress report to the Marine and Tropical Sciences Research Facility. Reef and Rainforest Research Centre Ltd, Cairns. 25pp.
- McKenzie, L.J., Lee Long, W.J., Coles, R.G. and Roder, C.A. (2000). Seagrass-Watch: Community based monitoring of seagrass resources. *Biol. Mar. Medit.* **7(2)**: 393-396.
- McKenzie, L.J., Campbell, S.J., Vidler, K.E. & Mellors, J.E. (2007) Seagrass-Watch: Manual for Mapping & Monitoring Seagrass Resources. Seagrass-Watch HQ, Cairns. 114pp
- Middlebrook, R., Williamson, J.E. (2006). Social attitudes towards marine resource management in two Fijian villages. *Ecological Management & Restoration* **7(2)**: 144-147.
- Smyth, D., Fitzpatrick, J., Kwan, D. (2006). Towards the development of cultural indicators for marine resource management in Torres Strait. CRC Torres Strait, Townsville. 61 pp.
- Turnbull, J. (2004). Explaining complexities of environmental management in developing countries: lessons from the Fiji Islands. *The Geographical Journal* **170** (1), 64-77.
- Waycott, M, McMahon, K, Mellors, J., Calladine, A., and Kleine, D (2004) A guide to tropical seagrasses in the Indo-West Pacific. (James Cook University Townsville) 72pp.

Further reading:

- Coles RG, McKenzie LJ and Campbell SJ. (2003). The seagrasses of eastern Australia. Chapter 11 In: World Atlas of Seagrasses. (EP Green and FT Short eds) Prepared by the UNEP World Conservation Monitoring Centre. (University of California Press, Berkeley. USA). pp 119-133.
- Costanza R, d'Arge R, de Groot R, Farber S, Grasso M, Hannon B, Limburg K, Naeem S, O'Neil RV, Paruelo J, Raskin RG, Sutton P and van der Belt M. (1997). The Value of the world's ecosystem services and natural capital. *Nature* **387**(15): 253-260.
- den Hartog C. (1970). The seagrasses of the world. (North-Holland Publishing, Amsterdam). 293pp.
- Green EP. and Short FT (Eds) (2003). World Atlas of Seagrasses. Prepared by the UNEP World Conservation Monitoring Centre. (University of California Press, Berkeley. USA). 298pp.
- Lanyon JM, Limpus CJ and Marsh H. (1989). Dugongs and turtles: grazers in the seagrass system. In: Biology of Seagrasses: A treatise on the biology of seagrasses with special reference to the Australian region. (AWD Larkum, AJ McComb and SA Shepherd eds). (Elsevier: Amsterdam, New York). pp 610-34.
- Larkum AWD, Orth RJ and Duarte CM (2006). Seagrasses: biology, ecology and conservation. Springer, The Netherlands. 691 pp.
- Orth RJ, Carruthers TJB, Dennison WC, Duarte CM, Fourqurean JW, Heck Jr KL, Hughes AR, Kendrick GA, Kenworthy WJ, Olyarnik S, Short FT, Waycott M and Williams SL. (2006). A Global Crisis for Seagrass Ecosystems. *BioScience* 56 (12): 987-996.
- Lee Long, W. J., Coles, R. G. & McKenzie, L. J. (2000) Issues for seagrass conservation management in Queensland. *Pacific Conservation Biology* 5, 321-328.
- Phillips, R.C, E.G Menez. (1988). Seagrasses. Smithsonian Institution Press, Washington, D.C. 104 pp.
- Poiner, I.R., Walker, D.I., and Coles, R.G. (1989). Regional Studies - Seagrass of Tropical Australia. In: Biology of Seagrasses. A.W.D. Larkum, A-J. McComb and S.A.Shepherd (Eds). Elsevier, Amsterdam, New York; 841 pp.
- Short, FT and Coles, RG. (Eds.) Global Seagrass Research Methods. Elsevier Science B.V., Amsterdam. 473pp.



Useful web links

Seagrass-Watch Official Site www.seagrasswatch.org

Western Australian Seagrass Webpage Focuses on Western Australian research and provides some general information and links to international seagrass sites.
www.science.murdoch.edu.au/centres/others/seagrass/

Seagrass Adventures Interactive website designed by students from Bentley Park College in Cairns (Australia). Website includes games, puzzles and quizzes for students to learn about seagrass and their importance. www.reef.crc.org.au/seagrass/index.html

World Seagrass Association A global network of scientists and coastal managers committed to research, protection and management of the world's seagrasses. wsa.seagrassonline.org

Seagrass Outreach Partnership Excellent website on seagrass of Florida. Provides some background information on seagrasses and has a great section with educational products and Seagrass Activity Kit for schools. www.flseagrass.org

Seagrass forum A global forum for the discussion of all aspects of seagrass biology and the ecology of seagrass ecosystems. Because of their complex nature, discussion on all aspects of seagrass ecosystems is encouraged, including: physiology, trophic ecology, taxonomy, pathology, geology and sedimentology, hydrodynamics, transplanting/restoration and human impacts. www.science.murdoch.edu.au/centres/others/seagrass/seagrass_forum.html

Reef Guardians and ReefEd Education site of the Great Barrier Reef Marine Park Authority. Includes a great collection of resources about the animals, plants, habitats and features of the Great Barrier Reef. Also includes an on-line encyclopedia, colour images and videos for educational use, a range of free teaching resources and activities. www.reefed.edu.au

Integration and Application Network (IAN) A website by scientists to inspire, manage and produce timely syntheses and assessments on key environmental issues, with a special emphasis on Chesapeake Bay and its watershed. Includes lots of helpful communication products such as fact sheets, posters and a great image library. ian.umces.edu

Reef Base A global database, information system and resource on coral reefs and coastal environments. Also extensive image library and online Geographic Information System (ReefGIS) which allows you to display coral reef and seagrass related data on interactive maps. www.reefbase.org

UNEP - World Conservation Monitoring Centre Explains the relationship between coral reefs, mangroves and seagrasses and contains world distribution maps. www.unep-wcmc.org

for more links, visit www.seagrasswatch.org/links.htm

We value your suggestions and any comments you may have to improve the Seagrass-Watch program.

Please complete the following statements in your own words

I found the Seagrass-Watch training to be

.....

.....

What I enjoyed most about the training was.....

.....

.....

It could have been better if.....

.....

.....

I did not realize that.....

.....

.....

Now I understand that.....

.....

.....

In my area the types of seagrasses and habitats include.....

.....

.....

.....

When I go back to my area, I will

.....
.....
.....

Other comments.....

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

Please hand in your form once you have completed it.



Thank you